




POLYOLEFIN PRODUCTION

Patent number: JP2002510358T
Publication date: 2002-04-02
Inventor:
Applicant:
Classification:
- international: C08F4/645; C08F10/00; C07F17/00
- european: C07F17/00; C08F10/00
Application number: JP19990512835T 19990219
Priority number(s): WO1999EP01094 19990219

Also published as:

 WO0049056 (A1)
 WO0049029 (A1)
 CA2371835 (A1)

Report a data error here

Abstract not available for JP2002510358T
Abstract of corresponding document: **WO0049029**

A metallocene catalyst component for use in preparing polyolefins having a monomer length of up to C10, which component has the general formula: $R''(CpR_1R_2R_3)(Cp'R_1'R_2')MQ_2$ wherein Cp is a substituted or unsubstituted cyclopentadienyl ring; Cp' is a substituted fluorenyl ring; R'' is a structural bridge imparting stereorigidity to the component; R1 is optionally a substituent on the cyclopentadienyl ring which is distal to the bridge, which distal substituent comprises a bulky group of the formula XR^*3 in which X is chosen from Group IVA, and each R* is the same or different and chosen from hydrogen or hydrocarbonyl of from 1 to 20 carbon atoms, R2 is optionally a substituent on the cyclopentadienyl ring which is proximal to the bridge and positioned non-vicinal to the distal substituent and is of the formula $YR\#3$ in which Y is chosen from group IVA, and each R# is the same or different and chosen from hydrogen or hydrocarbonyl of 1 to 7 carbon atoms, R3 is optionally a substituent on the cyclopentadienyl ring which is proximal to the bridge and is a hydrogen atom or is of the formula $ZR\text{ DOLLAR }3$, in which Z is chosen from group IVA, and each R DOLLAR is the same or different and chosen from hydrogen or hydrocarbonyl of 1 to 7 carbon atoms, R1' and R2' are each independently substituent groups on the fluorenyl ring, one of which is a group of the formula $AR'''3$, in which A is chosen from Group IVA, and each R''' is independently hydrogen or a hydrocarbonyl having 1 to 20 carbon atoms and the other is hydrogen or a second group of the formula $AR'''3$; M is a Group IVB transition metal or vanadium; and each Q is hydrocarbonyl having 1 to 20 carbon atoms or is a halogen.

Data supplied from the **esp@cenet** database - Worldwide